arXiv:1211.4505v1 [math.DS] 19 Nov 2012

STATISTICAL PROPERTIES OF QUADRATIC POLYNOMIALS WITH A NEUTRAL FIXED POINT

by

Artur Avila & Davoud Cheraghi

Abstract. — We describe the statistical properties of the dynamics of the maps $P_\alpha(z) := e^{2\pi\alpha\mathbf{i}}z + z^2$, with α of *high return* times. In particular, we show that these maps are *uniquely ergodic* on their *measure theoretic attractors*, and the unique invariant probability describes the statistical behavior of typical orbits in the *Julia set*. In particular, this confirms the conjecture of Perez-Marco on the unique ergodicity of *hedgehog* dynamics, in this class of maps.

Introduction

We study the statistical property of the orbits of the maps

$$P_\alpha(z) = e^{2\pi\alpha\mathbf{i}}z + z^2$$

on the complex plane, for irrational values of α.

The *post-critical set* of P_α is defined as

$$\mathcal{PC}(P_\alpha) := \overline{\cup_{j=1}^{\infty} P_\alpha^j(-e^{2\pi\alpha\mathbf{i}}/2)},$$

where $-e^{2\pi\alpha\mathbf{i}}/2$ is the unique critical point of the map. The geometry of this set and the dynamics of P_α on it play an important role in this study. A break point here is whether zero belongs to $\mathcal{PC}(P_\alpha)$ or not, which turns out to depend on the local dynamics of P_α at zero.

The map P_α is called *linearizable* at zero, if there exists a conformal change of coordinate ϕ defined near zero that fixes zero and conjugates P_α to the rotation of angle α about zero: $R_\alpha \circ \phi = \phi \circ P_\alpha$ on the domain of ϕ, where $R_\alpha(z) = e^{2\pi\alpha\mathbf{i}}z$. It was a nontrivial problem to determine the values of α for which P_α is linearizable. We refer the interested reader to [**Mil06**] for its rich history, and only need to mention that the answer to this problem depends on the arithmetic nature of α. For Lebesgue almost every $\alpha \in [0,1]$ such linearization exists, and on the other hand, for a generic choice of $\alpha \in [0,1]$, P_α is not linearizable. When P_α is linearizable at zero, the maximal domain on which P_α is conjugate to a rotation is called the *Siegel disk* of P_α.

2000 ***Mathematics Subject Classification.*** — Primary: 37F50, Secondary: 58F11.

By a result of Mane [**Mañ87**], the orbit of the critical point is recurrent, and when P_α is linearizable, the orbit of the critical point accumulates on the boundary of the Siegel disk, but when it is not linearizable, this orbit accumulates on zero. The inverse of the conjugacy ϕ from the disk of radius $1/2\pi$ to the Siegel disk of P_α provides us with an invariant probability measure defined on the boundary of the Siegel disk. That is, the push-forward of the Lebesgue measure on the boundary of the disk, which we denote by μ_α. The measure μ_α is the harmonic measure on the boundary of the Siegel disk, viewed from zero.

Our study uses a *renormalization* technique developed by H. Inou and M. Shishikura [**IS06**], and requires the entries in the continued fraction expansion of α be larger than a constant N, needed by this method. Define,

$$HT_N := \{[a_0, a_1, a_2, \dots,] \mid \forall i \geq 0, a_i \geq N\}.$$

Theorem A. — *There exists a constant $N > 0$ such that for $\alpha \in HT_N$, $P_\alpha : \mathcal{PC}(P_\alpha) \to \mathcal{PC}(P_\alpha)$ is uniquely ergodic. Moreover, when P_α is not linearizable at zero the unique invariant probability is the Dirac measure at 0, and when P_α is linearizable at zero, the unique invariant probability is μ_α.*

The above theorem and its proof have a number of applications to the statistical, geometric, and topological dynamics of these quadratics. According to a lemma of Lyubich [**Lyu83**], the post-critical set of a rational map, defined similarly as the closure of the forward orbits of its critical values, is the *measure theoretic attractor* of the dynamics of that map on its Julia set. This means that given any neighborhood of the post-critical set, the orbit of Lebesgue almost every point in the Julia set of the map eventually stays in that neighborhood. Combining this with the above theorem we obtain the following statistical property of the orbits of P_α.

Theorem B. — *For every $\alpha \in HT_N$ and Lebesgue almost every z in the Julia set of P_α, the measures*

$$\frac{1}{n} \sum_{j=0}^{n-1} \delta_{P_\alpha^j(z)}$$

converge to the unique invariant probability supported on $\mathcal{PC}(P_\alpha)$.

The theorem is meaningful only if the Julia set of P_α, $J(P_\alpha)$, has positive area. However, by a recent result of X. Buff and A. Chéritat [**BC12**], there are values of $\alpha \in [0,1]$ (of both types) where $J(P_\alpha)$ has positive area.(1) The above theorem applies to the examples by Buff and Chéritat, in particular.

Let f be a holomorphic germ defined on a neighborhood of zero with $f(0) = 0$ and $f'(0) = e^{2\pi\alpha\mathbf{i}}$ for an irrational α. Perez-Marco in [**PM97**] introduced local invariant sets for f, called *Siegel compacta*, that were used to study the local dynamics of f at zero. More precisely, he proves that for f as above and a neighborhood $U \ni 0$ where f is one-to-one on the closure of U, there exists a compact connected invariant set K with $0 \in K \subseteq \overline{U}$ and $K \cap \partial U \neq \emptyset$. A distortion property of the iterates of f on U is translated to the property of unique ergodicity of $f : \partial K \to \partial K$, which was conjectured to hold in this full generality. When f is not linearizable at zero, K has no interior ($\partial K = K$) and is called the *hedgehog* of f on U. It follows from Mane's [**Mañ93**] result that the boundary of every hedgehog of P_α must be contained in $\mathcal{PC}(P_\alpha)$. Hence, we can confirm the following partial result on this conjecture.

---

(1)It is expected that for a generic choice of $\alpha \in [0,1]$, $J(P_\alpha)$ has positive area.

Theorem C. — *For every $\alpha \in HT_N$ and every Siegel compacta K of P_α, the map $P_\alpha : \partial K \to \partial K$ is uniquely ergodic.*

On the topological side, note that Theorem A implies that there is no periodic point in the post-critical set, except possibly zero. As a byproduct of the proof of theorem A, we obtain an interesting property of the dynamics of linearizable maps counterpart to the *small cycle property* of nonlinearizable ones obtain by Yoccoz. He has shown that when P_α is not linearizable at zero, every neighborhood of zero contains infinitely many periodic cycles of P_α.

Theorem D. — *When P_α is linearizable at 0 for some $\alpha \in HT_N$, every neighborhood of the Siegel disk of P_α contains infinitely many periodic cycles of P_α.*

1. The renormalization of Inou and Shishikura

1.1. Inou-Shishikura class. — The cubic polynomial $P(z) := z(1+z)^2$ has a *parabolic* fixed point at 0, that is, a fixed point of multiplier $e^{2\pi\alpha \mathbf{i}}$ with $\alpha \in \mathbb{Q}$. Also, it has a critical point at $\mathrm{cp}_P := -1/3$ which is mapped to the critical value at $\mathrm{cv}_P := -4/27$, and another critical point at -1 which is mapped to zero.

Define the ellipse

$$E := \Big\{ x + \mathbf{i}y \in \mathbb{C} \mid (\frac{x+0.18}{1.24})^2 + (\frac{y}{1.04})^2 \leq 1 \Big\},$$

and consider

$$U := g(\hat{\mathbb{C}} \setminus E), \text{ where } g(z) := -\frac{4z}{(1+z)^2}. \tag{1}$$

The domain U contains 0 and cp_P, but not the other critical point of P at -1.

Inou and Shishikura in [**IS06**] define the class of maps

$$\mathcal{IS} := \Big\{ f := P \circ \varphi^{-1} \colon U_f \to \mathbb{C} \;\Big|\; \begin{array}{l} \varphi\colon U \to U_f \text{ is univalent, } \varphi(0) = 0,\ \varphi'(0) = 1, \\ \text{and } \varphi \text{ has a quasi-conformal extension to } \mathbb{C}. \end{array} \Big\}.$$

Every map in this class has a parabolic fixed point at 0 and a unique critical point at $\mathrm{cp}_f := \varphi(-1/3) \in U_f$. Given a set $A \subseteq \mathbb{R}$, we define

$$\mathcal{IS}_A := \{ f(e^{2\pi\alpha\mathbf{i}} z) \mid f \in \mathcal{IS}, \text{ and } \alpha \in A \}.$$

Abusing the notation, $\mathcal{IS}_\beta$ denotes the set $\mathcal{IS}_{\{\beta\}}$, for $\beta \in \mathbb{R}$.

There exists a natural Teichmüller metric on the class $\mathcal{IS}$. We do not work with this metric here, but only mention that the convergence in this metric implies the uniform convergence on compact sets, but not vice versa. Hence, by the sequence $h_n : \mathrm{Dom}\, h_n \to \mathbb{C}$, $n = 1, 2, \ldots$, *converges* to h we mean that given any compact set in $K \subseteq \mathrm{Dom}\, h$ and $\varepsilon > 0$, h_n is defined on K for sufficiently large n and uniformly tend to h on K. Note that the maps h_n are not necessarily defined on the same set.

The class $\mathcal{IS}_A$ naturally embeds into the space of univalent maps on the unit disk with a neutral fixed point at 0. Therefore, it is a precompact class in the compact-open topology.

Any map $h = e^{2\pi\alpha\mathbf{i}} f_0 \in \mathcal{IS}_\alpha$ has a fixed point at 0 with multiplier $e^{2\pi\alpha\mathbf{i}}$. Moreover, if α is small, h has another fixed point $\sigma_h \neq 0$ near 0 in U_h. The σ_h fixed point depends continuously on h and has asymptotic expansion $\sigma_h = -4\pi\alpha\mathbf{i}/f_0''(0) + o(\alpha)$, when h converges to f_0 in a fixed neighborhood of 0. Clearly $\sigma_h \to 0$ as $\alpha \to 0$.

We summarizes the basic local dynamics of maps in $\mathcal{IS}_\alpha$ proved by Inou and Shishikura. See Figure 1.

Theorem 1.1 **(Inou–Shishikura [IS06]).** — *There exist positive integers* $\boldsymbol{k}, \hat{\boldsymbol{k}}$, *and a real number* $\alpha_* > 0$ *satisfying the the following properties:*

1. *the set* $\{h''(0) \mid h \in \mathcal{IS}\}$ *is relatively compact in* $\mathbb{C} \setminus \{0\}$.
2. *For any map* $h\colon U_h \to \mathbb{C}$ *in* $\mathcal{IS}_\alpha$ *(or* $h = P_\alpha : \mathbb{C} \to \mathbb{C}$*) with* $\alpha \in (0, \alpha^*]$, *there exist a domain* $\mathcal{P}_h \subset U_h$ *and a univalent map* $\Phi_h\colon \mathcal{P}_h \to \mathbb{C}$ *satisfying the following properties:*
    - (a) *The domain* $\mathcal{P}_h$ *is bounded by piecewise smooth curves and is compactly contained in* U_h. *Moreover, it contains* cp_h, 0, *and* σ_h *on its boundary.*
    - (b) *There exists a continuous branch of argument defined on* $\mathcal{P}_h$ *such that*
    $$\max_{w,w' \in \mathcal{P}_h} |\arg(w) - \arg(w')| \leq 2\pi\hat{\boldsymbol{k}}.$$
    - (c) $\Phi_h(\mathcal{P}_h) = \{w \in \mathbb{C} \mid 0 < \mathrm{Re}(w) < \lfloor 1/\alpha \rfloor - \boldsymbol{k}\}$, $\mathrm{Im}\,\Phi_h(z) \to +\infty$ *when* $z \in \mathcal{P}_h \to 0$, *and* $\mathrm{Im}\,\Phi_h(z) \to -\infty$ *when* $z \in \mathcal{P}_h \to \sigma_h$.
    - (d) Φ_h *satisfies the Abel functional equation on* $\mathcal{P}_h$, *that is,*
    $$\Phi_h(h(z)) = \Phi_h(z) + 1, \text{ whenever } z \text{ and } h(z) \text{ belong to } \mathcal{P}_h.$$
    Furthermore, Φ_h *is unique once normalized by* $\Phi_h(\mathrm{cp}_h) = 0$.
    - (e) *The normalized map* Φ_h *depends continuously on* h.



FIGURE 1. A perturbed Fatou coordinate Φ_h and its domain of definition $\mathcal{P}_h$. The gray curve (amoeba) approximates the first few iterates of cp_h under h.

The map $\Phi_h : \mathcal{P}_h \to \mathbb{C}$ obtained in the above theorem is called the *perturbed Fatou coordinate*, or the *Fatou coordinate* for short, of h.

The class $\mathcal{IS}$ is denoted by $\mathcal{F}_1$ in [**IS06**]. All parts in the above theorem, except the existence of uniform $\hat{\boldsymbol{k}}$ and $\boldsymbol{k}$ in (b) and (c), follow readily from Theorem 2.1, Main Theorems 1, 3, and Corollary 4.2 in [**IS06**]. Parts (b) and (c) also follow from those results but require some extra work. A detailed proof of these statements are given in [**BC12**, Proposition 12].

Given a point z in the domain of a map $h : \operatorname{Dom} h \to \mathbb{C}$, the *orbit* of z under h, denoted by $\mathcal{O}(z)$, is defined as the sequence $(f^{\circ j}(z))_{j=0}^{m}$, where m is most number of times z can be iterated under h.

1.2. Renormalization. — Let $h\colon U_h \to \mathbb{C}$ either be in $\mathcal{IS}_\alpha$ or be P_α, with $\alpha \in (0, \alpha_*]$. Let $\Phi_h\colon \mathcal{P}_h \to \mathbb{C}$ denote the normalized Fatou coordinate of h. Define

$$\begin{aligned} \mathcal{C}_h &:= \{z \in \mathcal{P}_h : 1/2 \leq \operatorname{Re}(\Phi_h(z)) \leq 3/2\ ,\ -2 < \operatorname{Im}\Phi_h(z) \leq 2\}, \text{and} \\ \mathcal{C}_h^\sharp &:= \{z \in \mathcal{P}_h : 1/2 \leq \operatorname{Re}(\Phi_h(z)) \leq 3/2\ ,\ 2 \leq \operatorname{Im}\Phi_h(z)\}. \end{aligned} \tag{2}$$

By definition, $\mathrm{cv}_h \in \operatorname{int}(\mathcal{C}_h)$ and $0 \in \partial(\mathcal{C}_h^\sharp)$.

It follows from the work of Inou and Shishikura that there exists a positive integer k_h, depending on h, satisfying the following properties:

- For every integer k, with $0 \leq k \leq k_h$, there exists a unique connected component of $h^{-k}(\mathcal{C}_h^\sharp)$ which is compactly contained in $\operatorname{Dom} h$ and contains 0 on its boundary. We denote this component by $(\mathcal{C}_h^\sharp)^{-k}$.
- For every integer k, with $0 \leq k \leq k_h$, there exists a unique connected component of $h^{-k}(\mathcal{C}_h)$ which has non-empty intersection with $(\mathcal{C}_h^\sharp)^{-k}$, and is compactly contained in $\operatorname{Dom} h$. This component is denoted by $\mathcal{C}_h^{-k}$.
- The sets $\mathcal{C}_h^{-k_h}$ and $(\mathcal{C}_h^\sharp)^{-k_h}$ are contained in
$$\{z \in \mathcal{P}_h \mid \frac{1}{2} < \operatorname{Re}\Phi_h(z) < \frac{1}{\alpha} - \boldsymbol{k} - \frac{1}{2}\}.$$
- The maps $h : \mathcal{C}_h^{-k} \to \mathcal{C}_h^{-k+1}$, for $1 \leq k \leq k_h$, and $h : (\mathcal{C}_h^\sharp)^{-k} \to (\mathcal{C}_h^\sharp)^{-k+1}$, for $2 \leq k \leq k_h$, are univalent. The map $h : (\mathcal{C}_h^\sharp)^{-1} \to (\mathcal{C}_h^\sharp)$ is a degree two branched covering.

We define k_h as the smallest positive integer for which the above conditions hold, and let

$$S_h := \mathcal{C}_h^{-k_h} \cup (\mathcal{C}_h^\sharp)^{-k_h}.$$

By the Abel functional equation, the map

$$\Phi_h \circ h^{\circ k_h} \circ \Phi_h^{-1} : \Phi_h(S_h) \to \mathbb{C}. \tag{3}$$

projects via $z = \frac{-4}{27}e^{2\pi \mathbf{i} w}$ to a map $\mathcal{R}'(h)$ of the form $z \mapsto e^{2\pi \frac{-1}{\alpha}\mathbf{i}}z + O(z^2)$, defined on a set containing 0 in its interior. See Figure 2.

The conjugate map $s \circ \mathcal{R}'(h) \circ s^{-1}$, where $s(z) := \bar{z}$ denotes the complex conjugation map, has the form $z \mapsto e^{2\pi \frac{1}{\alpha}\mathbf{i}}z + O(z^2)$ near 0. The map $\mathcal{R}(h) := s \circ \mathcal{R}'(h) \circ s^{-1}$, restricted to the interior of $s(\frac{-4}{27}e^{2\pi \mathbf{i}(\Phi_h(S_h))})$, is called the *near-parabolic renormalization* of h by Inou and Shishikura. We simply



FIGURE 2. The figure shows the sets $\mathcal{C}_h$, $\mathcal{C}_h^\sharp$,..., $\mathcal{C}_h^{-k_h}$, and $(\mathcal{C}_h^\sharp)^{-k_h}$. The "induced map" projects via $e^{2\pi \mathbf{i} w}$ to a well defined map $\mathcal{R}(h)$ on a neighborhood of 0.

refer to it as the *renormalization* of h. For an slightly different definition of the same renormalization one may see [**BC12**].

We shall show in the next section (Lemma 2.1) that one time iterating $\mathcal{R}(h)$ corresponds to several times iterating h, via the change of coordinates. For some applications of a closely related renormalization (Douady-Ghys renormalization) one may see [**Dou87**], [**Dou94**], [**Yoc95**], [**Shi98**] and the references therein.

The following theorem of Inou and Shishikura [**IS06**, Main theorem 3] guarantees that the above definition of the near parabolic renormalization $\mathcal{R}$ can be carried out for small perturbations of maps in $\mathcal{IS}$. In particular, this implies the existence of k_h satisfying the three properties listed in the definition of the renormalization. See [**BC12**, Proposition 13] for a detailed argument on this. (2)

Define

$$V := P^{-1}(B(0, \frac{4}{27}e^{4\pi})) \setminus ((-\infty, -1] \cup B) \tag{4}$$

where B is the component of $P^{-1}(B(0, \frac{4}{27}e^{-4\pi}))$ containing -1 (see Figure 3).

---

(2) The sets $\mathcal{C}_h^{-k}$ and $(\mathcal{C}_h^\sharp)^{-k}$ defined here are (strictly) contained in the sets denoted by V^{-k} and W^{-k} in [**BC12**]. The set $\Phi_h(\mathcal{C}_h^{-k} \cup (\mathcal{C}_h^\sharp)^{-k})$ is contained in the union

$$D^\sharp_{-k} \cup D_{-k} \cup D''_{-k} \cup D'_{-k+1} \cup D_{-k+1} \cup D^\sharp_{-k+1}$$

in the notation used in [**IS06**, Section 5.A].



FIGURE 3. A schematic presentation of the polynomial P; its domain, and its range. Similar colors and line styles are mapped on one another.

By an explicit calculation (see [**IS06**, Proposition 5.2]) one can see that the closure of U is contained in the interior of V.

Theorem 1.2 **(Inou-Shishikura)**. — *There exist a constant* $\alpha^* > 0$ *such that if* $h \in \mathcal{IS}_\alpha$ *with* $\alpha \in (0, \alpha^*]$*, then* $\mathcal{R}(h)$ *is well-defined and belongs to the class* $\mathcal{IS}_{1/\alpha}$ *so that* $\mathcal{R}(h) := e^{\frac{2\pi}{\alpha}\boldsymbol{i}} \cdot P \circ \psi^{-1}$*. Moreover,* $\psi : U \to \mathbb{C}$ *extends to a univalent map on* V*.*

The same conclusion holds for the map $P_\alpha(z) = e^{2\pi\alpha\mathbf{i}}z + z^2$*. That is,* $\mathcal{R}(P_\alpha)$ *is well-defined and belongs to* $\mathcal{IS}_{1/\alpha}$ *provided* α *is small enough.*

Lemma 1.3. — *There exists a* $\boldsymbol{k}'' \in \mathbb{Z}$ *such that for every* $h \in \mathcal{IS}_{(0,\alpha_*]}$*,* $k_h \leq \boldsymbol{k}''$*.*

One may refer to [**Che10b**] for a proof of the above lemma. It is basically because $\mathcal{R}(h)$ extends over a larger domain definitely containing $\operatorname{Dom} \mathcal{R}(h)$ in its interior.

Let $[0; a_1, a_2, \dots]$ denote the continued fraction expansion of α as in the introduction. Define $\alpha_0 := \alpha$, and inductively for $i \geq 1$ define the sequence of real numbers $\alpha_i \in (0,1)$ as

$$\alpha_i := \frac{1}{\alpha_{i-1}} \pmod 1. \tag{5}$$

Then each α_i has expansion $[0; a_{i+1}, a_{i+2}, \dots]$.

If we fix a constant $N \geq 1/\alpha^*$, then $\alpha \in HT_N$ implies that $\alpha_j \in (0, \alpha^*)$, for $j = 0, 1, 2, \dots$. We use this constant N in the rest of this paper.

Let $\alpha \in HT_N$ and define $f_0 := P_\alpha$. Then, using Theorem 1.2, inductively define the sequence of maps

$$f_{n+1} := \mathcal{R}(f_n) : U_{f_{n+1}} \to \mathbb{C}.$$

Let $U_n := U_{f_n}$ denote the domain of definition of f_n, for $n \geq 0$. Hence, for every n,

$$f_n : U_n \to \mathbb{C},\ f_n(0) = 0, \text{ and } f_n'(0) = e^{2\pi\alpha_n\mathbf{i}}.$$

2. The renormalization tower

2.1. Changes of coordinates in the renormalization tower. —

Remark. — To slightly simplify the technical details of proofs, we assume that

$$N \geq \boldsymbol{k} + \hat{\boldsymbol{k}} + 1. \tag{6}$$

This is chosen so that $\Phi_{f_n}(\mathcal{P}_{f_n})$ is wide enough to contain a set defined later. However, this can be avoided by extending Φ_{f_n} and $\Phi_{f_n}^{-1}$ to larger domains, using the dynamics of f_n.

For $n \geq 0$, let $\Phi_n := \Phi_{f_n} : \mathcal{P}_n := \mathcal{P}_{f_n} \to \mathbb{C}$ denote the Fatou coordinate of $f_n \colon U_n \to \mathbb{C}$. We shall frequently use the notation for the map

$$\mathbb{E}\mathrm{xp}(\zeta) := \zeta \longmapsto \frac{-4}{27} s(e^{2\pi \mathbf{i} \zeta}) : \mathbb{C} \to \mathbb{C}^*, \text{ where } s(z) = \bar{z}.$$

By part (b) of Theorem 1.1, Inequality (6), and that $\mathcal{P}_n$ is simply connected, there is an inverse branch

$$\eta_n : \mathcal{P}_n \to \Phi_{n-1}(\mathcal{P}_{n-1})$$

of $\mathbb{E}\mathrm{xp}$. There may be several choices for this map but we choose one of them (for each n) such that

$$\mathrm{Re}(\eta_n(\mathcal{P}_n)) \subset [0, \hat{\boldsymbol{k}} + 1] \tag{7}$$

holds, and fix this choice for the rest of this note. Now define

$$\psi_n := \Phi_{n-1}^{-1} \circ \eta_n \colon \mathcal{P}_n \to \mathcal{P}_{n-1}.$$

Each ψ_n extends continuously to $0 \in \partial \mathcal{P}_n$ by mapping it to 0.

For $n \geq 2$ we can form the compositions

$$\Psi_n := \psi_1 \circ \psi_2 \circ \cdots \circ \psi_n \colon \mathcal{P}_n \to \mathcal{P}_0 \subset U_0.$$

For every $n \geq 0$, let $\mathcal{C}_n$ and $\mathcal{C}_n^\sharp$ denote the corresponding sets for f_n defined in (2) (i.e., replace h by f_n). Denote by k_n the smallest positive integer with

$$S_n^0 := \mathcal{C}_n^{-k_n} \cup (\mathcal{C}_n^\sharp)^{-k_n} \subset \{z \in \mathcal{P}_n \mid 0 < \mathrm{Re}\, \Phi_n(z) < \lfloor \frac{1}{\alpha_n} \rfloor - \boldsymbol{k} - 1\}.$$

By definition, the critical value of f_n is contained in $f_n^{\circ k_n}(S_n^0)$.

For every $n \geq 0$ and $i \geq 2$, define the sectors

$$S_n^1 := \psi_{n+1}(S_{n+1}^0) \subset \mathcal{P}_n, \text{and}$$
$$S_n^i := \psi_{n+1} \circ \cdots \circ \psi_{n+i}(S_{n+i}^0) \subset \mathcal{P}_n.$$

All these sectors contain 0 on their boundaries.

Lemma 2.1. — *Let $z \in \mathcal{P}_n$ be a point with $w := \mathbb{E}\mathrm{xp} \circ \Phi_n(z) \in U_{n+1}$. There exists an integer ℓ_z with $1 \leq \ell_z \leq \lfloor 1/\alpha_n \rfloor - \boldsymbol{k} + k_n - 1$, such that*

- *the orbit $z, f_n(z), f_n^{\circ 2}(z), \ldots, f_n^{\circ \ell_z}(z)$ is defined, $f_n^{\circ \ell_z}(z) \in \mathcal{P}_n$, and*
- $\mathbb{E}\mathrm{xp} \circ \Phi_n(f_n^{\circ \ell_z}(z)) = f_{n+1}(w)$

Proof. — As $w \in \operatorname{Dom} f_{n+1}$, by the definition of renormalization $\mathcal{R}(f_n) = f_{n+1}$, there are $\zeta \in \Phi_n(S_n^0)$ and $\zeta' \in \Phi_n(\mathcal{C}_n \cup \mathcal{C}_n^\sharp)$, such that

$$\mathbb{E}\mathrm{xp}(\zeta) = w, \quad \mathbb{E}\mathrm{xp}(\zeta') = f_{n+1}(w), \text{ and } \quad \zeta' = \Phi_n \circ f_n^{\circ k_n} \circ \Phi_n^{-1}(\zeta).$$

Since $\mathbb{E}\mathrm{xp}(\Phi_n(z)) = w$, there exists an integer ℓ with

$$-k_n + 1 \le \ell \le \lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1,$$

such that $\Phi_n(z) + \ell = \zeta$.

By the Abel functional equation for Φ_n, we have

$$\begin{aligned}\zeta' &= \Phi_n \circ f_n^{\circ k_n} \circ \Phi_n^{-1}(\zeta)\\ &= \Phi_n \circ f_n^{\circ k_n} \circ \Phi_n^{-1}(\Phi_n(z) + \ell)\\ &= \Phi_n \circ f_n^{\circ k_n + \ell}(z).\end{aligned}$$

Letting $\ell_z := k_n + \ell$, we have

$$1 \le \ell_z \le k_n + \lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1, \quad f_n^{\circ \ell_z}(z) = \Phi_n^{-1}(\zeta') \in \mathcal{P}_n, \text{ and}$$

$$\begin{aligned}\mathbb{E}\mathrm{xp} \circ \Phi_n(f_n^{\circ \ell_z}(z)) &= \mathbb{E}\mathrm{xp} \circ \Phi_n(\Phi_n^{-1}(\zeta'))\\ &= \mathbb{E}\mathrm{xp}(\zeta')\\ &= f_{n+1}(w).\end{aligned}$$

□

It should be clear that there are at least $\lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1$ choices for ℓ_z in the above lemma. In the following two lemmas we make some specific choices for ℓ_z to relate the number of iterates on level n to the number of iterates on level $n-1$, and also to the number of iterates on level 0. The proof is basically the uniqueness of the analytic continuation of conformal mappings that is presented in detail in [**Che10b**].

Define

$$\mathcal{P}_n' := \{w \in \mathcal{P}_n \mid 0 < \operatorname{Re} \Phi_n(w) < \lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1\}.$$

Lemma 2.2. — *For every $n \ge 1$ we have*

1. *for every $w \in \mathcal{P}_n'$, $f_{n-1}^{\circ \lfloor 1/\alpha_{n-1} \rfloor} \circ \psi_n(w) = \psi_n \circ f_n(w)$, and*
2. *for every $w \in S_n^0$, $f_{n-1}^{\circ (k_n \lfloor 1/\alpha_{n-1} \rfloor + 1)} \circ \psi_n(w) = \psi_n \circ f_n^{\circ k_n}(w)$.*

Lemma 2.3. — *For every $n \ge 1$ we have*

1. *for every $w \in \mathcal{P}_n'$, $f_0^{\circ q_n} \circ \Psi_n(w) = \Psi_n \circ f_n(w)$,*
2. *for every $w \in S_n^0$, $f_0^{\circ (k_n q_n + q_{n-1})} \circ \Psi_n(w) = \Psi_n \circ f_n^{\circ k_n}(w)$, and*
3. *similarly, for every $m < n$, $f_n \colon \mathcal{P}_n' \to \mathcal{P}_n$ and $f_n^{\circ k_n} : S_n^0 \to (\mathcal{C}_n \cup \mathcal{C}_n^\sharp)$ are conjugate to some iterates of f_m on the set $\psi_{m+1} \circ \cdots \circ \psi_n(\mathcal{P}_n)$.*

2.2. The approximating neighborhoods. — For every $n \geq 0$, consider the union

$$\Omega_n^0 := \bigcup_{i=0}^{k_n+\lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1} f_n^{\circ i}(S_n^0) \bigcup \{0\} \subset \operatorname{Dom} f_n. \tag{8}$$

Using Lemma 2.3, we transfer the iterates in the above union to the dynamic plane of f_0 to define

$$\Omega_0^n := \bigcup_{i=0}^{q_n(k_n+\lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1)+q_{n-1}} f_0^{\circ i}(S_0^n) \bigcup \{0\}.$$

The upper bound in the above union is obtained as follows. The first k_n iterates in (8) correspond to $k_n q_n + q_{n-1}$ number of iterates on level 0 by Lemma 2.3–(2), and the remaining $\lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1$ iterates in (8) amounts to $q_n(\lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1)$ number of iterates by Lemma 2.3–(1). The main result of this section is the following proposition whose proof can be found in [**Che10b**].

Proposition 2.4. — *For every $n \geq 0$, we have*

(1) *Ω_0^{n+1} is compactly contained in the interior of Ω_0^n;*

(2) *$\mathcal{PC}(f_0)$ is contained in the interior of Ω_0^n.*

It follows from [**Lyu83**] that the orbit of almost every point in the Julia set of P_α eventually stays in each Ω^n. That is, for every $n \geq 0$ and almost every $z \in J(P_\alpha)$ there is a positive integer k with $\mathcal{O}(f^{\circ k}(z) \subset \Omega^n$.

The next lemma (see [**Che10b**] for the proof) proves that while the orbit of a point stays in some Ω^n, it visits all the sectors involved in that union.

Lemma 2.5. — *$\forall n \geq 0$, $\forall \ell \in \mathbb{Z}$ with $0 \leq \ell \leq q_n(k_n + \lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1) + q_{n-1}$, and for almost every $z \in J(f_0)$, $\mathcal{O}(z) \cap f_0^{\circ \ell}(S_0^n) \neq \emptyset$.*

3. Unique ergodicity of post-critical dynamics

We will use the following equivalent definition of unique ergodicity, see [**Mañ87**, Theorem 9.2]. A continuous map $T : X \to X$ of a compact metric space X is uniquely ergodic if for every continuous function $f : X \to \mathbb{R}$ and every $x \in X$ the limit

$$\lim_{n\to\infty} \frac{1}{n} \sum_{j=0}^{n-1} f(T^{\circ j}(x))$$

exists and is independent of x.

For every $n \in \mathbb{N}$, let

$$l_n := q_n(k_n + \lfloor 1/\alpha_n \rfloor - \boldsymbol{k} - 1) + q_{n-1},$$
$$s_n := \min_{z \in \mathcal{PC}(P_\alpha) \cap S_0^n} \min\{k \in \{2,3,4,\dots\} \mid P_\alpha^{\circ k}(z) \in S_0^n\}.$$

By Proposition 2.4, $s_n \leq l_n$. For every $\delta > 0$ and $n \in \mathbb{N}$ define the set

$$C(n,\delta) := \left\{ j \in \{0,1,2,\dots,l_n\} \;\middle|\; P_\alpha^{\circ j}(S_0^n) \subseteq B(0,\delta) \right\}.$$

Proposition 3.1. — *For every non-Brjuno $\alpha \in HT_N$ and every $\delta > 0$, we have*

$$\limsup_{n\to\infty} \frac{|C(n,\delta)| + s_n}{l_n} = 2.$$

For Brjuno values of α let Δ_α denote the Siegel disk of P_α, and $B_\delta(\Delta_\alpha)$ the δ neighborhood this disk. Given $n \in \mathbb{N}$ and $\delta > 0$, define $S(n,\delta)$ as the maximal element (by cardinality) of the set

$$\left\{ \{a, a+1, a+2, \dots, b\} \;\middle|\; \begin{array}{l} 0 \le a \le b \le l_n \text{ and} \\ \forall k \in \{a, a+1, \dots, b\}, P_\alpha^{\circ k}(S_0^n) \subseteq B_\delta(\Delta_\alpha) \end{array} \right\}.$$

Proposition 3.2. — *For every Brjuno $\alpha \in HT_N$ we have the following.*

(1)

$$\lim_{\substack{\delta\to 0\\ n\to\infty}} \max_{k\in S(n,\delta)} \operatorname{diam}\,(P_\alpha^{\circ k}(S_0^n) \setminus \Delta_\alpha) = 0.$$

(2) *For every $\delta > 0$*

$$\limsup_{n\to\infty} \frac{|S(n,\delta)| + s_n}{l_n} = 2.$$

Assuming these two propositions, which are proved in the following two sections, we give a proof of Theorem A. Although both of linearizable and non-linearizable maps can be treated simultaneously. As the proof in the non-linearizable case is slightly easier and demonstrates the idea for the other case, we present them separately.

Proof of Theorem A for non-Brjuno values of α. — Let $f : \mathcal{PC}(P_\alpha) \to \mathbb{R}$ be a continuous map with supremum norm $M := \|f\|_\infty$. Let ε be an arbitrary positive constant and choose $\delta' > 0$ such that for all $w \in B(0,\delta')$ we have $|f(w) - f(0)| < \varepsilon$.

Fix a non-zero point $z \in \mathcal{PC}(P_\alpha)$ and $n \in \mathbb{N}$. Let $t_0, t_1, t_2, \cdots$ denote the consecutive moments of the returns of the orbit of z to S_0^n. That is, t_0 is the smallest positive integer with $P_\alpha^{\circ t_0}(z) \in S_0^n$. Then assuming we have t_i, t_{i+1} is the smallest integer larger that $t_i + 1$ with $P_\alpha^{\circ t_{i+1}}(z) \in S_0^n$. By definition,

$$|t_0| \le l_n, \text{ and } \forall i \ge 0, s_n \le |t_{i+1} - t_i| \le l_n + 1.$$

Given $N \in \mathbb{N}$ large enough, choose $m \in \mathbb{N}$ with $t_m < N \leq t_{m+1}$.

$$\begin{aligned}
&\Big|\frac{1}{N}\sum_{k=0}^{N-1} f(P_\alpha^{\circ k}(z)) - f(0)\Big| \\
&\qquad = \Big|\frac{1}{N}\Big(\sum_{k=0}^{t_0-1} f(P_\alpha^{\circ k}(z)) + \sum_{k=t_m}^{N-1} f(P_\alpha^{\circ k}(z)) + \sum_{i=1}^{m}\sum_{k=t_{i-1}}^{t_i-1} f(P_\alpha^{\circ k}(z))\Big) - f(0)\Big| \\
&\qquad \leq \frac{1}{N}\Big(l_n 2M + l_n 2M + \sum_{i=1}^{m}\sum_{k=t_{i-1}}^{t_i-1} \big|f(P_\alpha^{\circ k}(z)) - f(0)\big|\Big) \\
&\qquad \leq \frac{1}{N}\Big(l_n 4M + \sum_{i=1}^{m}\big(C(n,\delta)\varepsilon + (l_n - C(n,\delta))2M\big)\Big) \\
&\qquad \leq \frac{1}{N}\Big(l_n 4M + \frac{N}{s_n}\big(C(n,\delta)\varepsilon + (l_n - C(n,\delta))2M\big)\Big).
\end{aligned}$$

As $N \to \infty$ the last expression tends to

$$\frac{C(n,\delta)}{s_n}\varepsilon + \frac{l_n - C(n,\delta)}{s_n}2M$$

If $n \to \infty$ along the sequence obtained in Proposition 3.1, the above quantity tends to ε. Therefore,

$$\lim_{N\to\infty} \frac{1}{N}\sum_{k=0}^{N-1} f(P_\alpha^{\circ k}(z)) = f(0)$$

□

Proof of Theorem A for Brjuno values of α. — Let $f : \mathcal{PC}(P_\alpha) \to \mathbb{R}$ be continuous with $M := \|f\|_\infty$. Let ε be an arbitrary positive constant and choose $\delta' > 0$ such that for all z, w in $\mathcal{PC}(P_\alpha)$ with $|z - w| < \delta'$ we have $|f(w) - f(z)| < \varepsilon$. Now, using Proposition 3.2 choose $n \in \mathbb{N}$ and $\delta > 0$ for which we have

$$\max_{k\in S(n,\delta)} \operatorname{diam}(P_\alpha^{\circ k}(S_0^n) \setminus \Delta_\alpha) < \delta'.$$

Fix arbitrary points $z \in \mathcal{PC}(P_\alpha)$ and $z' \in \partial\Delta_\alpha$. Let t_i, for $i = 0, 1, 2, \cdots$, (respectively, t'_i, for $i = 0, 1, 2, \cdots$) be the moments of the visits of the orbit of z (respectively, z') to S_0^n. That is, t_0 is the smallest positive integer with $P_\alpha^{\circ t_0}(z) \in S_0^n$. Assuming t_i is defined, t_{i+1} is the smallest positive integer larger than $t_i + 1$ with $P_\alpha^{\circ t_{i+1}}(z) \in S_0^n$. The sequence t'_i is defined similarly for the orbit of z'.

We have

$$|t_0| \leq l_n, \text{ and } \forall i \geq 0, s_n \leq |t_{i+1} - t_i| \leq l_n + 1,$$

as well as

$$|t'_0| \leq l_n, \text{ and } \forall i \geq 0, s_n \leq |t'_{i+1} - t'_i| \leq l_n + 1.$$

Given a large enough integer N, choose $m' \in \mathbb{N}$ with $t'_{m'} < N \leq t'_{m'+1}$, as well as $m \in \mathbb{N}$ with $t_m < N \leq t_{m+1}$. Note that m, m' are bounded by $\frac{N}{s_n}$, and $|m - m'|$ is bounded by $\frac{N}{s_n} - \frac{N}{l_n+1}$.

Let $B_n := \cup_{j\in S(n,\delta)} P_\alpha^{\circ j}(S_0^n)$. For the average along the orbit of z' we have

$$\begin{aligned}
&\Big|\frac{1}{N}\sum_{k=0}^{N-1} f(P_\alpha^{\circ k}(z')) - \frac{1}{N}\sum_{i=1}^{m'} \sum_{\substack{k=t'_{i-1}\\ P_\alpha^{\circ k}(z')\in B_n}}^{t'_i-1} f(P_\alpha^{\circ k}(z'))\Big| \\
&\qquad = \frac{1}{N}\Big|\sum_{k=0}^{t'_0-1} f(P_\alpha^{\circ k}(z')) + \sum_{k=t'_{m'}}^{N-1} f(P_\alpha^{\circ k}(z')) + \sum_{i=1}^{m'} \sum_{\substack{k=t'_{i-1}\\ P_\alpha^{\circ k}(z')\notin B_n}}^{t'_i-1} f(P_\alpha^{\circ k}(z'))\Big| \\
&\qquad \le \frac{1}{N}\Big(2l_n + \frac{N}{s_n}(l_n - S(n,\delta))\Big)M
\end{aligned}$$

Similarly, for the average along the orbit of z we obtain

$$\begin{aligned}
&\Big|\frac{1}{N}\sum_{k=0}^{N-1} f(P_\alpha^{\circ k}(z)) - \frac{1}{N}\sum_{i=1}^{m} \sum_{\substack{k=t_{i-1}\\ P_\alpha^{\circ k}(z)\in B_n}}^{t_i-1} f(P_\alpha^{\circ k}(z))\Big| \\
&\qquad \le \frac{1}{N}\Big(2l_n + \frac{N}{s_n}(l_n - S(n,\delta))\Big)M
\end{aligned}$$

On the other hand,

$$\begin{aligned}
&\Big|\frac{1}{N}\sum_{i=1}^{m'} \sum_{\substack{k=t'_{i-1}\\ P_\alpha^{\circ k}(z')\in B_n}}^{t'_i-1} f(P_\alpha^{\circ k}(z')) - \frac{1}{N}\sum_{i=1}^{m} \sum_{\substack{k=t_{i-1}\\ P_\alpha^{\circ k}(z)\in B_n}}^{t_i-1} f(P_\alpha^{\circ k}(z))\Big| \\
&\le \frac{1}{N}\Big|\sum_{i=1}^{\min\{m,m'\}} \Big(\sum_{\substack{k=t'_{i-1}\\ P_\alpha^{\circ k}(z')\in B_n}}^{t'_i-1} f(P_\alpha^{\circ k}(z')) - \sum_{\substack{k=t_{i-1}\\ P_\alpha^{\circ k}(z)\in B_n}}^{t_i-1} f(P_\alpha^{\circ k}(z))\Big) + \sum_{i=\min\{m,m'\}}^{\max\{m,m'\}} \sum_{\substack{k=r_{i-1}\\ P_\alpha^{\circ k}(w)\in B_n}}^{r_i-1} M\Big| \\
&\le \frac{1}{N}\left(\frac{N}{s_n}\,|S(n,\delta)|\,\varepsilon + (\frac{N}{s_n} - \frac{N}{l_n+1})M\right) \\
&\le \frac{S(n,\delta)}{s_n}\,\varepsilon + (\frac{1}{s_n} - \frac{1}{l_n+1})M
\end{aligned}$$

Putting the above three estimates together, we conclude

$$\begin{aligned}
&\Big|\frac{1}{N}\sum_{k=0}^{N} f(P_\alpha^{\circ k}(z')) - \frac{1}{N}\sum_{k=0}^{N} f(P_\alpha^{\circ k}(z))\Big| \\
&\qquad \le \frac{4l_nM}{N} + \frac{l_n - S(n,\delta)}{s_n}2M + \frac{S(n,\delta)}{s_n}\,\varepsilon + (\frac{1}{s_n} - \frac{1}{l_n+1})M.
\end{aligned}$$

Letting $N \to \infty$, and then $n \to \infty$ along the sequence provided by Proposition 3.2-(2), and that ε is arbitrarily, we come to

$$\lim_{N\to\infty} \frac{1}{N} \sum_{k=0}^{N-1} f(P_\alpha^{\circ k}(z)) = \lim_{N\to\infty} \frac{1}{N} \sum_{k=0}^{N-1} f(P_\alpha^{\circ k}(z')) = \int_{\partial\Delta_\alpha} f \, d\mu.$$

□

Proof of Corollary B. — It follows from [**Lyu83**] (or see [**Che10b**, Proposition 1.2]) that the limit set of Lebesgue almost every point in $J(P_\alpha)$ is contained in $\mathcal{PC}(P_\alpha)$. (Indeed, in [**Che10a**] we show that this limit set is equal to $\mathcal{PC}(P_\alpha)$.) For any such point z, every convergent subsequence of the sequence of measures

$$\mu_n := \frac{1}{n} \sum_{k=0}^{n-1} \delta_{P_\alpha^{\circ k}(z)}.$$

is invariant under P_α and is supported on $\mathcal{PC}(P_\alpha)$. However, by Theorem A there is only one invariant measure supported on $\mathcal{PC}(P_\alpha)$. Hence, the above sequence of measures is convergent and converges either to Dirac measure at 0 or μ, depending on the type of α. □

4. Proof of Propositions 3.1 and 3.2

The proof consists of two ingredients. An arithmetic one that has been analysed by Yoccoz in [**Yoc95**], and an analytic part that has been developed in [**Che10a**].

4.1. An arithmetic lemma.

— Given an irrational number α, recall the sequence of numbers $(\alpha_i)_{i=0}^\infty$ defined by Equation (5). Let $\beta_0 := 1$, and $\beta_k := \Pi_{i=1}^k \alpha_i$, $k \geq 1$.

Let B be an arbitrary real constant and define the sequence of numbers

$$\begin{aligned} &B_{k,k} := -2, \qquad k = 0, 1, 2, \cdots, \\ &B_{k,i-1} := \alpha_i B_{k,i} + \log \alpha_i^{-1} - B, \qquad 1 \leq i < k. \end{aligned} \tag{9}$$

Lemma 4.1. — *For every irrational α and every $B \in \mathbb{R}$ we have,*

1. *For every $k \geq 0$*

$$B_{k,0} \leq \sum_{i=1}^{k} \beta_{i-1} \log \alpha_i^{-1} \leq B_{k,0} + 3B + 2,$$

2. *If α is non-Brjuno, for every $T \in \mathbb{R}$ the set*

$$\mathcal{L}(\alpha, T) := \{k \in \mathbb{N} \mid B_{k,i} \geq T\alpha_i^{-1}, \textit{for } 1 < i < k\}$$

has infinite cardinality.

(3) *For $T \in \mathbb{R}$ and $m \in \mathbb{N}$ let*

$$\mathcal{L}(\alpha, T, l) := \{k \in \{l+1, l+2, \dots\} \mid B_{k,i} \geq T\alpha_i^{-1}, \text{ for } l < i < k\}$$

Assume that α is a Brjuno number and for some T, $\mathcal{L}(\alpha, T, l)$ has finite cardinality for every $l \in \mathbb{N}$, then

$$\liminf_{j\to\infty} \lim_{m\to\infty} B_{m,j} - T\alpha_j^{-1} < \infty.$$

The first two statements of the above lemma, when one starts with 0 instead of -2 in Equation (9) and $B = T$, were proved in [**Yoc95**, Section 1.6]. The same argument works here as well, but for the sake of completeness we present a proof here.

Proof. — *Part (1):* Recursively using (9), we have

$$\begin{aligned}
&B_{k,i-1}\\
&\quad= \alpha_i B_{k,i} + \log \alpha_i^{-1} - B\\
&\quad= \alpha_i(\alpha_{i+1} B_{k,i+1} + \log \alpha_{i+1}^{-1} - B) + \log \alpha_i^{-1} - B\\
&\quad= \alpha_i\alpha_{i+1} B_{k,i+1} + \alpha_i \log \alpha_{i+1}^{-1} + \log \alpha_i^{-1} - B(1+\alpha_i)\\
&\quad\ \ \vdots\\
&\quad= \beta_{i-1}^{-1}\big(\beta_k B_{k,k} + \beta_{k-1}\log\alpha_k^{-1} + \beta_{k-2}\log\alpha_{k-1}^{-1} + \cdots + \beta_{i-1}\log\alpha_i^{-1} - B(\beta_{i-1} + \beta_i + \cdots + \beta_{k-1})\big)\\
&\quad= -2\beta_{i-1}^{-1}\beta_k + \beta_{i-1}^{-1}\sum_{j=i}^{k}\beta_{j-1}(\log\alpha_j^{-1} - B).
\end{aligned}$$

In particular, for $i = 1$, using the inequality $\alpha_j\alpha_{j+1} < 1/2$, we conclude the first part of the lemma.

Part (2): Assume on the contrary that $\mathcal{L}(\alpha, T)$ is finite for some α. Let k denote the largest element in $\mathcal{L}(\alpha, T) \cup \{0\}$.

Given a positive integer $N_0 > k$, define the integers $N_0 > N_1 > N_2 > \cdots > N_r \leq k$ according to

$$B_{N_{l-1}, N_l} < T\alpha_{N_l}^{-1}, \quad \text{for } l = 1, 2, \dots, r.$$

By the equation in the proof of Part (1), this implies that

$$\sum_{N_l+1}^{N_{l-1}} \beta_{j-1}(\log\alpha_j^{-1} - B) \leq 2\beta_{N_{l-1}} + T\beta_{N_l-1}.$$

Adding these sums together, one obtains

$$\sum_{j=N_r+1}^{N_0} \beta_{j-1}\log\alpha_j^{-1} \leq 2\sum_{l=1}^{r}\beta_{N_{l-1}} + T\sum_{l=1}^{r}\beta_{N_l-1} + B\sum_{j=N_r+1}^{N_0}\beta_{j-1}.$$

As $N_0 \to \infty$,

$$\sum_{j=k}^{\infty} \beta_{j-1} \log \alpha_j^{-1} \leq 2(2+T+B),$$

contradicting our assumption on the type of α.

Part (3): By definition each set $\mathcal{L}(\alpha, T, l)$ contains $l+1$, and therefore is non-empty. Define $\ell_0 := 0$, and $\ell_{n+1} := \max \mathcal{L}(\alpha, T, \ell_n)$, for $n \in \mathbb{N}$. This implies that

$$B_{\ell_{n+1}, \ell_n} < T\alpha_{\ell_n}^{-1}, \quad \text{for } n = 0, 1, 2, \dots.$$

Now fix $k \in \mathbb{N}$, and let $n > k$ be a arbitrary integer.

For every $j \in \{k+1, k+2, \dots, n\}$ we have

$$|B_{\ell_j, \ell_{j-1}} - 2| < T\alpha_{\ell_{j-1}}^{-1} + 2,$$

which by Definition (9) implies

$$\begin{aligned} |B_{\ell_j, \ell_k} - B_{\ell_{j-1}, \ell_k}| &< (T\alpha_{\ell_{j-1}}^{-1} + 2)\alpha_{\ell_{j-1}}\alpha_{\ell_{j-1}-1} \dots \alpha_{\ell_k+1} \\ &= T\beta_{\ell_{j-1}-1}\beta_{\ell_k}^{-1} + 2\beta_{\ell_{j-1}}\beta_{\ell_k}^{-1}. \end{aligned}$$

Putting all these together for $j = k+1, k+2, \dots, n$ we obtain

$$\begin{aligned} |B_{\ell_n, \ell_k} - 2| &\leq \sum_{j=k+1}^{n} T\beta_{\ell_{j-1}-1}\beta_{\ell_k}^{-1} + 2\beta_{\ell_{j-1}}\beta_{\ell_k}^{-1} \\ &\leq T\alpha_{\ell_k}^{-1} + 2(T+2). \end{aligned}$$

If an integer $m \notin \{\ell_k, \ell_{k+1}, \ell_{k+2}, \dots\}$, choose n with $\ell_{n-1} < m < \ell_n$. By definition of the sequence ℓ_n we have

$$B_{\ell_n, m} \geq T\alpha_m^{-1} > B_{m,m}.$$

This implies that

$$B_{\ell_n, \ell_k} > B_{m, \ell_k}.$$

All in all, we have shown that

$$\sup_m B_{m, \ell_k} \leq T\alpha_{\ell_k}^{-1} + 2(T+2).$$

which implies the wanted statement. □

4.2. The heights of chains in the tower. — Given natural numbers $n > m$, and a point $\zeta_n \in \operatorname{Dom} \chi_n$, let

$$\mathcal{C}_{n,m}(\zeta_n) := \left\{ \langle \zeta_n, \zeta_{n-1}, \dots, \zeta_m \rangle \;\middle|\; \begin{array}{l} \zeta_j \in \operatorname{Dom} \chi_j, \ \Phi_j \circ \mathbb{E}\mathrm{xp}(\zeta_{j-1}) = \zeta_j \\ \text{for } j = n, n-1, \dots, m-1 \end{array} \right\}.$$

An element $\tau := \langle \zeta_n, \zeta_{n-1}, \dots, \zeta_m \rangle$ in the above set may be referred to as a *chain of length* $n-m+1$ *starting at* ζ_n, and also, for every $k \in \{n, n-1, \dots, m\}$, we define $\tau_k := \zeta_k$. The plan is to study the behaviour of the numbers $\langle \tau_j \rangle_{j=n}^{0}$, for $\tau \in \mathcal{C}_{n,0}(t)$, $t \in \operatorname{Dom} \chi_n$. This depends on the arithmetic of α, and requires some fine estimates on the (Perturbed) Fatou coordinates.

Let σ_j denote the non-zero fixed point of f_j (that is near 0). The change of coordinate Φ_j^{-1} can be decomposed as $T_j \circ L_j^{-1}$, where T_j is the covering map from $\mathbb{C}$ to $\hat{\mathbb{C}} \setminus \{0, \sigma_j\}$ defined as

$$T_j := \frac{\sigma_j}{1 - e^{-2\pi\alpha_j w \mathbf{i}}}.$$

The map L_j may be defined as $\Phi_j \circ T_j$. Through this decomposition, one lifts f_j via T_j, to obtained a map F_j that can be estimated using basic Taylor or Köebe estimates on f_j. The map L_j conjugates F_j to the translation by 1, and can be estimated (through non trivial processes) from estimates on F_j. One may refer to [**Yoc95, Shi98, Shi00, Che10b, Che10a**], among other works, for further details on this. In particular in [**Che10a**] we prove the following estimate.

For every $n \in \mathbb{N}$ let $\chi_n : \Phi_n(\mathrm{Dom}\, \Phi_n) :\rightarrow \mathbb{C}$ denote an inverse branch of $\Phi_n^{-1} \circ \mathbb{E}\mathrm{xp}$.

Proposition 4.2. — *$\forall D > 0$, $\exists M$ such that $\forall j \in \mathbb{N}$ and $\forall w \in \mathrm{Dom}\, \chi_j$ with $\mathrm{Im}\, w \geq D/\alpha_j$,*

$$|\chi_j'(w) - \alpha_j| \leq M \alpha_j e^{-2\pi\alpha_j \,\mathrm{Im}\, w}.$$

Moreover, as $D \to \infty$, $M \to 1$.

Note that the above estimate is independent of the choice of the lift in χ_j.

Let $B_{n,j}$ denote the sequences of points defined in the previous subsection. Define, $\Delta_j' := \Phi_j(\Delta_j)$, for $j \in \mathbb{N}$.

Lemma 4.3. — *$\exists B, M \in \mathbb{R}$ such that $\forall \alpha \in HT_N$ and $\forall n \in \mathbb{N}$, there exists a $\tau \in \mathcal{C}_{n,0}(\frac{1}{2\alpha_n} - 2\mathbf{i})$ such that*

$$B_{n,j} \leq \mathrm{Im}\, \tau_j \leq B_{n,j} + M, \text{ for } j = 0, 1, 2, \ldots, n-1.$$

(Here, the sequence $B_{k,i}$ is defined with the constant B.)

Moreover,

$$\sup_{j \in \mathbb{N}} \Big| \lim_{n \to \infty} B_{n,j} - \max_{w \in \partial \Delta_j'} \mathrm{Im}\, w \Big| < \infty.$$

Proof. — For every n, we choose $\tau := \langle \frac{1}{2\alpha_n} - 2\mathbf{i}, \zeta_{n-1}, \ldots, \zeta_0 \rangle \in \mathcal{C}_{n,0}(\frac{1}{2\alpha_n} - 2\mathbf{i})$ according to the rule $\mathrm{Re}\, \zeta_j \in [\frac{1}{2\alpha_j} - \frac{1}{2}, \frac{1}{2\alpha_j} + \frac{1}{2})$. To prove the lemma we need to introduce a constant B (independent of n and j) such that

$$\mathrm{Im}\, \tau_{j-1} \geq \alpha_j \,\mathrm{Im}\, \tau_j + \log \frac{1}{\alpha_j} - B, \text{ for } j = n, n-1, \ldots, 1. \tag{10}$$

Given $\tau_j \in \mathrm{Dom}\, \chi_j$, with $\mathrm{Re}\, \zeta_j \in [\frac{1}{2\alpha_j} - \frac{1}{2}, \frac{1}{2\alpha_j} + \frac{1}{2})$, it follows from Lemmas 5.2 and 5.4 in [**Che10b**] that $L_j^{(}\tau_j)$ belongs to the disk of radius $C - \log \alpha_j$ centered at τ_j, for some constant C independent of n and j. Then, it is an explicit calculation to show that Equation (10) holds for some constant B independent of n and j. One may refer to the proof of Proposition 3.3 in [**Che10b**] for further details. ☐

Using Proposition 4.2, choose $D > 0$ such that for every $\alpha \in (0,1)$ the following inequalities hold.

$$\int_{D/\alpha}^{\infty} M\alpha e^{-2\pi\alpha t}\, dt \leq 1/2, \text{ and } \int_{-\boldsymbol{k}''}^{1/\alpha+\boldsymbol{k}''} M\alpha e^{-2\pi\alpha D}\, dt \leq 3/2. \tag{11}$$

Lemma 4.4. — *With the constant D obtained above we have the following. If for some $n \in \mathbb{N}$ there exists a $\tau \in \mathcal{C}_{n,0}(\frac{1}{2\alpha_n} - 2\mathbf{i})$ satisfying*

$$\operatorname{Im} \tau_j \geq \frac{D+12}{\alpha_j}, \text{ for } j = 0, 1, 2, \dots, n-1,$$

then for every $\tau' \in \mathcal{C}_{n,0}(\frac{1}{2\alpha_n} - 2\mathbf{i})$ we have

$$\operatorname{Im} \tau'_j \geq \operatorname{Im} \tau_j - 12, \text{ for } j = 0, 1, 2, \dots, n-1.$$

Proof. — Note that

$$D + 4(1 + \alpha_1 + \alpha_1\alpha_2 + \alpha_1\alpha_2\alpha_3 + \alpha_1\alpha_2\alpha_3\alpha_4 + \dots) \leq D + 12,$$

since $\alpha_j\alpha_{j+1} < 1/2$, for all j.

Assume $\tau \in \mathcal{C}_{n,0}(\frac{1}{2\alpha_n} - 2\mathbf{i})$ be a chain satisfying the hypotheses of the lemma, and $\tau' \in \mathcal{C}_{n,0}(\frac{1}{2\alpha_n} - 2\mathbf{i})$ be an arbitrary chain. For every $j = n-1, n-2, \dots, 1$, let γ_j denote the piecewise horizontal and vertical curve connecting τ_j to τ'_j. If the curve γ_j is above the horizontal line passing through $D\mathbf{i}/\alpha_j$, we have

$$\begin{aligned} |\operatorname{Im}(\tau_{j+1} - \tau'_{j+1})| &= |\operatorname{Im} \int_{\gamma_j} \chi'_j\, dz| \\ &\leq \operatorname{Im} \int_{\gamma_j} |\chi'_j - \alpha_j|\, dz + |\operatorname{Im} \int_{\gamma_j} \alpha_j\, dz| \\ &\leq 4 + \alpha_j |\operatorname{Im}(\tau_j - \tau'_j)|. \end{aligned}$$

One infers the conclusion from an inductive argument. □

4.3. Geometry of sectors on level n.

— For every $n \geq 1$, let R_n denote the maximal element (by its cardinality) of the set

$$\left\{ \{a, a+1, \dots, b\} \;\middle|\; \begin{array}{l} a, b \in \{0, 1, 2, \dots, \lfloor 1/\alpha_n \rfloor - \boldsymbol{k}\}, \text{ and for } a \leq k \leq b \\ \inf \operatorname{Im} \mathbb{L}og(f_n^k(S_n^0)) \geq \frac{1}{2} \operatorname{Im} \chi_n(\frac{1}{2\alpha_n} - 2\mathbf{i}) \end{array} \right\}.$$

If the above set is empty, we let $R_n := \emptyset$.

Lemma 4.5. — $\forall \varepsilon > 0$, $\exists \delta > 0$ *such that if $\alpha_n < \delta$, then*

$$1 - \frac{|R_n|}{\lfloor 1/\alpha_n \rfloor + \boldsymbol{k}_n - \boldsymbol{k}} < \varepsilon. \tag{12}$$

Proof. — Recall the digits $[a_0, a_1, a_2, \dots,]$ of the continued fraction expansion of α_0. We claim that for small enough α_n, the infinite strip

$$B_n := \left\{ w \in \mathbb{C} \mid \operatorname{Im} w \in [-2, \infty), \operatorname{Re} w \in [a_n^{2/3}, a_n - a_n^{2/3}] \right\},$$

lifts under χ_n to a set above the height $2^{-1} \log 1/\alpha_n$. This implies the lemma.

Recall the map L_n that conjugates the lift F_n to the translation by one. To prove the above claim, it is enough to show that for sufficiently small α_n we have

$$L_n^{-1}(B_n) \subseteq \hat{B}_n := \left\{ w \in \mathbb{C} \mid \operatorname{Im} w \in [\log \alpha_n, \infty), \operatorname{Re} w \in [a_n^{2/3}/2, a_n - a_n^{2/3}/2] \right\}.$$

This is because by an explicit calculation of the maps T_n and $\mathbb{E}\mathrm{xp}$, $\hat{B}_n$ is mapped to a set above the height $\frac{-2}{3} \log \alpha_n - C$, for some constant C independent of n.

To locate the set $L_n^{-1}(B_n)$, we need some elementary estimates on L_n and F_n from [**Che10b**]. Let $\gamma_n(t) := a_n/2 + t\mathbf{i}, t \geq -2$ (a ray at the center of B_n), which is mapped to $\gamma'_n := L_n^{-1} \circ \gamma_n$. By Lemma 5.2-4, $\gamma'_n(0)$ lies within the ball of radius $-C \log \alpha_n$ centered at $a_n/2$, for some constant C independent of n. Also, by Lemma 5.4, the curve γ'_n lies within a uniformly bounded distance, say C, from a vertical line. To obtain the image of B_n under L_n^{-1}, we need to iterate F_n on γ'_n, forward and backward, $(a_n/2 - a_n^{2/3})$ number of times. As F_n is nearly a translation by one, with explicit bounds in Lemma 5.2-2, these iterates reach at most the vertical lines passing through $a_n^{2/3} + 2C \log \alpha_n - C$ and $a_n - a_n^{2/3} - C \log \alpha_n + C$.

On the other hand, by the same discussion, $|\operatorname{Im} \chi_n(\frac{1}{2\alpha_n} - 2\mathbf{i}) + \log \alpha_n|$ is uniformly bounded by a constant independent of n. Hence, for small enough α_n, this set is contained in $\hat{B}_n$. ☐

We also need the following easy lemma.

Lemma 4.6. — *If a subsequence $\langle \alpha_{n_k} \rangle_{k=0}^{\infty}$ of the sequence $\langle \alpha_n \rangle_{n=0}^{\infty}$ converges to zero as $k \to \infty$, then*

$$\lim_{k \to \infty} \frac{s_{n_k}}{l_{n_k}} = 1.$$

Now we are ready to prove the following stronger statement that implies Proposition 3.1.

Proposition 4.7. — *For every non-Brjuno $\alpha \in HT_N$ there exists a sequence of real numbers δ_i, $i = 0, 1, 2 \dots$, converging to zero such that*

$$\limsup_{n \to \infty} \frac{|C(n, \delta_n)| + s_n}{l_n} = 2.$$

Proof. — Let T_n, $n = 1, 2, 3, \cdots$, be an increasing sequence of real numbers all bigger than $2D + 48$. Define the increasing sequence of positive integers $n_i := \min \mathcal{L}(\alpha, T_i)$, for $i = 1, 2, 3, \dots$, obtained in Lemma 4.1-2. It follows that the sequence $\alpha_{n_i} \to 0$, as $i \to \infty$. Hence by Lemma 4.6, we have

$$\lim_{i \to \infty} \frac{s_{n_i}}{l_{n_i}} = 1.$$

Recall the sequence $B_{n_i,0}$, $i \geq 1$, defined in Equation (9), and choose $\delta_i \in \mathbb{R}$, $i \in \mathbb{N}$, such that

$$\Phi_0^{-1}\left\{w \in \operatorname{Dom}\Phi_0^{-1} \mid \operatorname{Im} w \geq B_{n_i,0} - \frac{1}{2}\alpha_1\alpha_2\dots\alpha_{n_i-1}\log\frac{1}{\alpha_{n_i}} - 24\right\} \subseteq B_{\delta_i}(0).$$

From Lemma 4.1-1, and that α_0 is a non-Brjuno number, one concludes that $\delta_i \to 0$ as $i \to \infty$.

By virtue of Lemma 4.5,

$$\lim_{i\to\infty}\frac{|R_{n_i}|}{\lfloor 1/\alpha_{n_i}\rfloor + \boldsymbol{k}_{n_i} - \boldsymbol{k}} = 1.$$

We claim that all the lifts of the sectors

$$f_{n_i}^k(S_{n_i}^0), k \in R_{n_i}$$

to the first level of renormalization provide us with sectors contained in $B_{\delta_i}(0)$. Assuming this for a moment, from the definition of the renormalization, the domains Ω^n, and Lemmas 2.3 and 2.2 we conclude

$$P_\alpha^{\circ j}(S_0^n) \subseteq B(0,\delta_i), \text{ for } q_{n_i-1} + \min R_{n_i} q_{n_i} \leq j \leq q_{n_i-1} + \max R_{n_i} q_{n_i}.$$

Hence, the percentage of the sectors in Ω^{n_i} that fit in $B_{\delta_i}(0)$ tends to one.

To prove the claim fix an integer $n = n_i$ in the above sequence. We need to prove that for every $w \in B_n$ such that all its consecutive lifts $w_n := w, w_{n-1}, w_{n-2}, \dots, w_0$ under χ_j, respectively, are defined, then

$$\operatorname{Im} w_0 \geq B_{n,0} - \frac{1}{2}\alpha_1\alpha_2\dots\alpha_n\log\frac{1}{\alpha_n} - 24.$$

By Lemma 4.3, there exists $\tau \in \mathcal{C}_{n,0}(\frac{1}{2\alpha_n} - 2\mathbf{i})$ with

$$\operatorname{Im}\tau_j \geq B_{n,j} \geq \frac{2D+48}{\alpha_j}, \text{ for } j = n-1, n-2, \dots, 0.$$

Therefore, by Lemma 4.4, for $\tau' \in \mathcal{C}_{n,0}(\frac{1}{2\alpha_n} - 2\mathbf{i})$ that is defined along the same branches as the sequence $\langle w_j\rangle$, we must have

$$\operatorname{Im}\tau'_j \geq B_{n,j} - 12, \text{ for } j = 0,1,2,\dots,n-1. \tag{13}$$

By an inductive procedure we show that for all $j = 0,1,2,\dots,n-1$

$$|\operatorname{Im}(\tau'_j - w_j)| \leq \frac{1}{2}\alpha_{j+1}\alpha_{j+2}\dots\alpha_{n-1}\log\frac{1}{\alpha_n} + 4(1 + \alpha_{j+1} + \alpha_{j+1}\alpha_{j+2} + \dots). \tag{14}$$

For $j = n - 1$ the inequality follows from the definition of the set B_n. Now assume we have Equation (14) for some j. Combining with Equation (13), $\operatorname{Im} w_j$ enjoys

$$\begin{aligned}\operatorname{Im} w_j &\geq \operatorname{Im} \tau'_j - |\operatorname{Im} \tau'_j - \operatorname{Im} w_j| \\ &\geq B_{n,j} - 12 - (\frac{1}{2}\alpha_{j+1}\alpha_{j+2}\dots\alpha_{n-1}\log\frac{1}{\alpha_n} + 4(1+\alpha_{j+1}+\alpha_{j+1}\alpha_{j+2}+\dots) \\ &\geq B_{n,j} - \frac{1}{2}\alpha_{j+1}\alpha_{j+2}\dots\alpha_{n-1}\log\frac{1}{\alpha_n} - 24 \\ &\geq \frac{1}{2}B_{n,j} - 24 \geq \frac{D}{\alpha_j}\end{aligned}$$

Now choose a piecewise horizontal and vertical curve γ_j connecting w_j to τ'_j that lies above the height D/α_j. Since w_j and τ'_j belong to the image of the same inverse branch, their real part is at most k' apart. The estimate in the proof of Lemma 4.4 implies Equation (14) for $j + 1$.

Now Equations 13 and 14 for $j = 0$ imply the desired estimate on the height of w_0. ☐

4.4. The linearizable maps. — We consider the following two scenarios. Let Δ_m, for $m = 0, 1, 2, \dots$, denote the Siegel disk of the m-th renormalization map f_m.

$\mathfrak{A}$ **:** For every $T \in \mathbb{R}$ there exist $m_0 \in \mathbb{N}$ and infinitely many integers $m > m_0$ such that $\mathcal{C}_{m,0}(\frac{1}{2\alpha_m} - 2\mathbf{i})$ contains an element τ with

$$\operatorname{Im} \tau_j > \frac{T}{\alpha_j}, \text{ for } j = m-1, m-2, \dots, m_0 + 1.$$

$\mathfrak{B}$ **:** There exist constants T and M, as well as infinitely many levels m such that

$$\forall w \in \operatorname{Dom} \chi_m, \text{with } \operatorname{Im} w \geq \frac{T}{\alpha_m} + M,\ \Phi_m^{-1}(w) \in \Delta_m.$$

Lemma 4.8. — *For every irrational rotation α, at least one of the above statements holds.*

Proof. — If $\mathfrak{A}$ does not hold for some T, then it does not hold for larger values of T as well. Using Lemma 4.3, there exists a constant T' such that $\mathcal{L}(\alpha, T', l)$ is finite for every $l \in \mathbb{N}$, where $\mathcal{L}(\alpha, T', l)$ is defined in Lemma 4.1-3. Hence, by Lemmas 4.1-3, there exists a sequence of levels $\ell_0, \ell_1, \ell_2, \dots$ on which

$$\lim_{m\to\infty} B_{m,\ell_j} - T'\alpha_{\ell_j}^{-1} < \infty.$$

Using Lemma 4.3, we conclude that for every $m \in \mathbb{N}$, and every $\tau \in \mathcal{C}_{m,0}(\frac{1}{2\alpha_m} - 2\mathbf{i})$ we have

$$\operatorname{Im} \tau_{\ell_j} - T\alpha_{\ell_j}^{-1} < \infty.$$

Since the interior of $\cap_{m\geq 0}\Omega_{\ell_j}^m$ contains the Siegel disk of f_{ℓ_j}, we obtain the statement $\mathfrak{B}$. ☐

Lemma 4.9. — *The derivatives of the changes of coordinates from level n to level 0 tends to 0, as n tends to infinity.*

Proof. — The proof depends on how the lifts are defined. It follows from a hyperbolic contraction, or, the properties of the map on the Siegel disk, and definition of the Julia set. ☐

Proof of Proposition 3.2 when $\mathfrak{A}$ holds. — The proof is similar to the one for nonlinearizable maps.

By Lemma 4.3,

$$H := \sup_{j \in \mathbb{N}} \big| \lim_{n \to \infty} B_{n,j} - \max_{w \in \partial \Delta'_j} \operatorname{Im} w \big| + 25,$$

is a finite number.

By the assumption for the constant $2D + 48$, there exists a level m_0, and levels $m_1, m_2, \dots$ such that each $\mathcal{C}_{m_i,0}(\frac{1}{2\alpha_{m_i}} - 2\mathbf{i})$ contains an element τ with

$$\operatorname{Im} \tau_j > \frac{2D + 48}{\alpha_j}, \text{ for } j = m_i - 1, m_i - 2, \dots, m_0 + 1.$$

Fix $\delta \in (0, \infty)$. By virtue of Lemma 4.9, we can replace m_0 by a larger integer so that for every $m > m_0$, every $w \in B_H(\Delta_m)$, and every $\tau \in \mathcal{C}_{m,0}(w)$, we have $\Phi_0^{-1}(\tau_0) \in B_\delta(\Delta_0)$.

Recall the set R_{m_i} defined in Section 4.3. We want to show that for sufficiently large m_i, all the lifts of the sectors

$$f_{m_i}^k(S_{m_i}^0), k \in R_{m_i}$$

to level m_0 provide us with sectors contained in $B_H(\Delta'_{m_0})$. This is because by the estimate in the proof of Proposition 4.7, for every $w \in B_{m_i}$, and every $\tau' \in \mathcal{C}_{m_i,0}(w)$ we have

$$\operatorname{Im} \tau_{m_0} \geq B_{m_i,m_0} - \frac{1}{2} \alpha_{m_0+1} \alpha_{m_0+2} \dots \alpha_{m_i-1} \log \frac{1}{\alpha_{m_i}} - 24.$$

Moreover, since the Brjuno sum for α_{m_0} is finite, the second term on the right hand side of the above inequality is eventually less than one.

The sequence $\langle \alpha_{m_i} \rangle_{i=1}^{\infty}$ must tend to zero because of the following reason. For every $\tau \in \mathcal{C}_{m_i,0}$, $\operatorname{Im} \tau_{m_i-1}$ is nearly $-\log \alpha_{m_i}$. Now if $\log \alpha_{m_i}$ is uniformly bounded independent of i, then by the contraction in Lemma 4.9 and that the point one lifts to the point one, the critical value must belong to the boundary of the Siegel disk. Contradicting our assumption. Hence, we have

$$\lim_{i \to \infty} \frac{|R_{n_i}|}{\lfloor 1/\alpha_{n_i} \rfloor + \boldsymbol{k}_{n_i} - \boldsymbol{k}} = 1.$$

This finishes the proof of the Proposition. ☐

Proof of Proposition 3.2 when $\mathfrak{B}$ holds. — Let the sequence m_i, $i = 1, 2, 3, \dots$ denotes the levels on which the inequality in $\mathfrak{B}$ holds. Consider the sets

$$B_{m_i,j} := \{w \in \mathbb{C} \mid \operatorname{Im} w \geq -2, j \leq \operatorname{Re} w \leq j + 1, \Phi_{m_i}^{-1}(w) \notin \Delta_{m_i}\}, \text{for } j = 0, 1, 2, \dots, \lfloor \alpha_{m_i}^{-1} \rfloor - \boldsymbol{k} - 1.$$

We need to show that for every δ, most of these sets, when lifted to the level zero, are contained in $B_\delta(\Phi_0(\Delta_0))$.

If the sequence α_{m_i}, has a subsequence bounded away from zero, then on those levels the sets $B_{m_i,j}$, for all j, have uniformly bounded diameter. Therefore, by Lemma 4.9, the diameters of all the lifts to the top level tend to zero, and hence the conclusion of the proposition follows.

Now assume that $\alpha_{m_i} \to 0$ as $i \to \infty$ and fix a constant $\delta > 0$. Recall the constant $\hat{\boldsymbol{k}}$ chosen such that the lift of $\Phi_n(\operatorname{Dom}\Phi_n)$ to the previous level horizontally spans at most diameter $\hat{\boldsymbol{k}}$. In particular, for large enough i, we have the inequalities

$$\frac{1}{2\alpha_{m_i}} + 2\hat{k} \leq \frac{1}{\alpha_{m_i}} - \boldsymbol{k}, \text{ and } \frac{1}{2\alpha_{m_i}} + 2\hat{k} \leq \frac{2}{3\alpha_{m_i}}.$$

Consider the sets

$$C_n := \{w \in \mathbb{C} \mid \operatorname{Im} w \geq -2, \frac{1}{4\alpha_n} - \hat{\boldsymbol{k}} \leq \operatorname{Re} w \leq \frac{3}{4\alpha_n} + \hat{\boldsymbol{k}}\}.$$

By the estimate in Lemma 4.2, the lifts $\chi_n(C_n)$ have uniformly bounded diameter M independent of n. Moreover, by Lemma 4.9, there exists an integer m' such that every set of diameter M on a level larger than m', is sent to a set of diameter at most δ on level 0 under the lifts. Also, note that from all the lifts $\chi_n(\operatorname{Dom}\chi_n)$ that are contained in the domain of χ_{n-1}, at list half of them are contained in C_{n-1}. This implies that as m_i tends to infinity, the ratio of the lifts of the sets $B_{m_i,j}$ that lie within $B_\alpha(\Phi_0(\Delta_0))$ tends to one. ☐

Remark. — It will be fruitful to find a condition similar to the statement $\mathfrak{A}$ that is equivalent to the critical value being off the boundary of the Siegel disk of P_α.

Proof of Theorem D. — This basically follows from the above proofs of Proposition 3.2. More precisely, when $\mathfrak{A}$ occurs, the σ_m fixed point of f_m lifts under $\mathbb{E}\mathrm{xp}$ to a set of points with imaginary more than $\log\alpha_m^{-1}$ minus a uniform constant. By the definition of the renormalization this set of points are mapped under Φ_{m-1} to a periodic point for f_{m-1}. By an inductive process and the argument in the proof of Proposition 3.2 in this case, we obtain a cycle with in small neighborhood of the Siegel disk.

Now assume m is one of the levels on which the inequality in $\mathfrak{B}$ has occurred. By the estimate in Proposition 4.2, there are points on the boundary of the Siegel disk that are uniformly bounded away from the lifts of the σ_m-fixed point of f_m. Then by the contraction in Lemma 4.9, these lifts of σ_m lift to a periodic point for P_0 near the Siegel disk Δ_0. ☐

Proof of Corollary C. — We use a theorem of Perez-Marco and a theorem of Mane, to show that the boundary of every hedgehog of P_α is contained in $\mathcal{PC}(P_\alpha)$. Indeed, Given a hedgehog K of P_α, by [**PM95**], there exists a sequence of iterates $P_\alpha^{n_i}$ that converges to the identity on K. On the other hand, by [**Mañ93**] on any subset of the Julia set that avoids the post-critical set of P_α, the backward iterates of P_α are contracting.

☐

References

[BC12] X. Buff & A. Chéritat – "Quadratic Julia sets with positive area", *Ann. of Math. (2)* **176** (2012), no. 2, p. 673–746.

[Che10a] D. Cheraghi – "Typical orbits of quadratic polynomials with a neutral fixed point", To appear in Comm. Math. Phy., preprint 2012, 2010.

[Che10b] ———, "Typical orbits of quadratic polynomials with a neutral fixed point: non-Brjuno type", Preprint, 2010.

[Dou87] A. Douady – "Disques de Siegel et anneaux de Herman", *Astérisque* (1987), no. 152-153, p. 4, 151–172 (1988), Séminaire Bourbaki, Vol. 1986/87.

[Dou94] ———, "Does a Julia set depend continuously on the polynomial?", in *Complex dynamical systems (Cincinnati, OH, 1994)*, Proc. Sympos. Appl. Math., vol. 49, Amer. Math. Soc., Providence, RI, 1994, p. 91–138.

[IS06] H. Inou & M. Shishikura – "The renormalization for parabolic fixed points and their perturbation", Preprint, 2006.

[Lyu83] M. Lyubich – "Typical behavior of trajectories of the rational mapping of a sphere", *Dokl. Akad. Nauk SSSR* **268** (1983), no. 1, p. 29–32.

[Mañ87] R. Mañé – *Ergodic theory and differentiable dynamics*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 8, Springer-Verlag, Berlin, 1987, Translated from the Portuguese by Silvio Levy.

[Mañ93] ———, "On a theorem of Fatou", *Bol. Soc. Brasil. Mat. (N.S.)* **24** (1993), no. 1, p. 1–11.

[Mil06] J. Milnor – *Dynamics in one complex variable*, third ed., Annals of Mathematics Studies, vol. 160, Princeton University Press, Princeton, NJ, 2006.

[PM95] R. Pérez Marco – "Sur une question de Dulac et Fatou", *C. R. Acad. Sci. Paris Sér. I Math.* **321** (1995), no. 8, p. 1045–1048.

[PM97] R. Pérez-Marco – "Fixed points and circle maps", *Acta Math.* **179** (1997), no. 2, p. 243–294.

[Shi98] M. Shishikura – "The Hausdorff dimension of the boundary of the Mandelbrot set and Julia sets", *Ann. of Math. (2)* **147** (1998), no. 2, p. 225–267.

[Shi00] ———, "Bifurcation of parabolic fixed points", in *The Mandelbrot set, theme and variations*, London Math. Soc. Lecture Note Ser., vol. 274, Cambridge Univ. Press, Cambridge, 2000, p. 325–363.

[Yoc95] J.-C. Yoccoz – "Théorème de Siegel, nombres de Bruno et polynômes quadratiques", *Astérisque* (1995), no. 231, p. 3–88, Petits diviseurs en dimension 1.

November 20, 2012

Artur Avila • *E-mail :* `artur@math.sunysb.edu`, CNRS UMR 7586, Institut de Mathe matiques de Jussieu, 175 rue du Chevaleret, 75013-Paris, FRANCE

Davoud Cheraghi • *E-mail :* `d.cheraghi@warwick.ac.uk`, Mathematics Institute, University of Warwick, Coventry, CV4 7AL, Engalnd